

Novel use of lumen-apposing metal stent for recanalization of complete stenosis of colo-colonic anastomosis

OPEN
ACCESS

Combined antegrade-retrograde dilation [1] is an established technique that has been adapted for benign colonic obstruction [2,3] but requires use of accessory devices and predilation followed by deployment of self-expanding metal stents [2,3]. Rendezvous sigmoidorectal reanastomosis using a lumen-apposing metal stent (LAMS) has also been described with endosonographic (EUS) guidance [4]. We hereby report a recanalization of complete stenosis of a colo-colonic anastomosis without EUS guidance.

An 81-year-old woman with a history of large-bowel obstruction secondary to benign sigmoid diverticular stricture underwent a two-step diverting transverse loop colostomy creation followed by transverse loop colostomy takedown with colo-colonic anastomosis, open sigmoidectomy with colorectal anastomosis, and diverting loop ileostomy creation. Subsequent barium enema and sigmoidoscopy showed a patent colorectal anastomosis but complete stenosis of the colo-colonic anastomosis in the distal transverse colon (► Fig. 1). Given the distant location of the stenosis from either orifice, we opted for combined antegrade-retrograde forward-viewing colonoscopy for recanalization of the colon lumen.

Using fluoroscopy and transillumination, a 15 × 15 mm LAMS (AXIOS; Boston Scientific, Marlborough, Massachusetts, USA) was safely deployed across the stenosis (► Fig. 2a, b). Balloon dilation was performed within the saddle of the stent (► Fig. 2c).

The patient was discharged home and a subsequent colonoscopy was performed 2 months later for retrieval of the LAMS and dilation of the remnant anastomosis to 18 mm (► Video 1). Given patent colo-colonic and colo-rectal anastomosis, the patient underwent loop ileostomy takedown and had complete remission of symptoms with regular bowel movements.



► **Fig. 1** Initial findings. **a** Barium enema with contrast filling terminating at the distal transverse colon. **b** Schematic diagram demonstrating anatomy of the patient's colon. Source for ► Fig. 1 b: Lance Powell.




► **Fig. 2** Recanalization procedure. **a** Combined antegrade-retrograde positioning of endoscopes on fluoroscopy. **b** Lumen-apposing metal stent (LAMS) puncture across the anastomosis. **c** Dilation of the LAMS post deployment.

LAMS allows puncture and release in a single-step procedure, thus enabling deployment of the stent into the target lumen without prior guidewire insertion or preliminary dilation, thereby simplifying the procedure with low enteroanastomosis migration risk [5]. Although EUS guidance is typically utilized, this may be technically challenging or unavailable; however, safe LAMS deployment can be performed using only transillumination and fluoroscopy.

Endoscopy_UCTN_Code_TTT_1AQ_2AF

Acknowledgments

The authors would like to thank Lance Powell and Rebecca Harrison for artistic rendering of the patient's anatomy and assistance with formatting the images, respectively.



Video 1 Retrieval of the lumen-apposing metal stent and dilation of the colo-colonic anastomosis.

Bibliography

Endoscopy 2023; 55: E470–E471

DOI 10.1055/a-2013-1820

ISSN 0013-726X

© 2023. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany



ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



Endoscopy E-Videos is an open access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online. Processing charges apply, discounts and waivers acc. to HINARI are available.

This section has its own submission website at <https://mc.manuscriptcentral.com/e-videos>

Competing interests

The authors declare that they have no conflict of interest.

The authors

Paul Guzik¹, Allison Aka², Wasseem Skef¹

- 1 Division of Gastroenterology and Hepatology, Loma Linda University Medical Center, Loma Linda, California, United States
- 2 Division of Colon and Rectal Surgery, Loma Linda University Medical Center, Loma Linda, California, United States

Corresponding author

Wasseem Skef, MD

Division of Gastroenterology and Hepatology, Loma Linda University Medical Center, 11234 Anderson Street, Loma Linda, CA 92354, United States
wassem.skef@gmail.com

References

- [1] van Twisk JJ, Brummer RJ, Manni JJ. Retrograde approach to pharyngo-esophageal obstruction. *Gastrointest Endosc* 1998; 48: 296–299
- [2] Moyer MT, Mathew A, Chintanaboina J et al. Restoration of colonic patency of a completely obstructed Crohn's stricture using the combined antegrade-retrograde dilation procedure. *VideoGIE* 2017; 2: 359–360
- [3] Ngamruengphong S, Mohapatra S, Runge T et al. Recanalization of a complete coloanal anastomotic obstruction using a combined antegrade-retrograde rendezvous technique. *Endoscopy* 2020; 52: E312–E314
- [4] Kozieł S, Kozłowska-Petriczko K, Pawlak K et al. Endoscopic sigmoidorectal reanastomosis using a dual endoscope technique: rendezvous single-balloon enteroscopy and endoscopic ultrasound. *Endoscopy* 2021; 53: E257–E258
- [5] Garcia-Alonso FJ, Sanchez-Ocana R, Peñas-Herrero I et al. Cumulative risks of stent migration and gastrointestinal bleeding in patients with lumen-apposing metal stents. *Endoscopy* 2018; 50: 386–395